

# Features of Kamiokande-II, IMB and Baksan observations and their interpretation in a two-component model for the signal

Francesco Vissani<sup>1</sup> and Giulia Pagliaroli<sup>2</sup>

<sup>1</sup>*INFN, Laboratori Nazionali del Gran Sasso, Assergi (AQ), Italy*

<sup>2</sup>*University of L'Aquila, Coppito, and INFN, Laboratori Nazionali del Gran Sasso, Italy*

We consider the time, angular and energy distributions of SN1987A events discussing the quality of their agreement with the expectations. A global interpretation is performed considering a simple model, based on the standard scenario for the explosion. Despite the contrasting and confusing indications, a straightforward fit to the data provides a result that does not contradict but rather supports the expectations. The calculated electron antineutrino flux is applied to predict the relic neutrino signal.

## I. INTRODUCTION

The Symposium held in Moscow on February 2007 [1] was devoted to discuss open problems of SN1987A. The meaning of LSD events [2] was discussed, and the potential of the model of Imshennik and Ryazhskaya [3] was shown. The observations of Kamiokande-II [4], IMB [5] and Baksan [6] were covered by the contribution of Alexeev. In the spirit of the Symposium, we recall that (1) certain features of these observations are difficult to understand, see e.g. [7]; (2) there is not yet a complete model for the explosion, that could shed doubts on the assumption that we know how neutrinos are emitted; this applies also to ref. [3].

In this work that expands the talk of F.V. at [1] we study the features of Kamiokande-II, IMB and Baksan data and analyze them in the ‘standard scenario’ of emission, first advocated by Nadyozhin [8]—also called Bethe and Wilson scenario [9], delayed explosion or  $\nu$ -assisted explosion. We recall its features: On top of a several-seconds lasting emission from the proto neutron star, the **cooling**, there is an initial intense emission lasting a fraction of a second. The  $\nu_e$  and  $\bar{\nu}_e$  irradiated in this phase transfer energy and help the stalled shock wave to resume, eventually leading to explosion. We call this phase of neutrino emission **accretion**. For progresses toward the implementation of the ‘standard scenario’ in numerical simulations, see [10].

## II. FEATURES OF THE DATA

The observed events are 11 or 12 in Kamiokande-II, the 6<sup>th</sup> being below the threshold of 7.5 MeV. The duration and the energy of the signal is not safely known. Thus we consider a unified time window of  $T = 30$  s in this work. The candidate signal events increases to 16 in Kamiokande-II (the events number 13,14,15,16 being below the threshold), it is 8 in IMB and 5 in Baksan. Recall that most signal is due to  $\bar{\nu}_e p \rightarrow e^+ n$  [IBD reaction] that has the largest cross section; unless stated otherwise we assume that all signal events are due to IBD.

The background in Kamiokande-II declines rapidly with the number of hit phototubes; on average, there

should be about 0.7 background events above threshold and 5.6 in total. Kamiokande-II analyzed the whole volume of the detector. The region close to the walls (in particular the upper wall) is the less safe against the risk of background. Five of the 11 events are close to the walls, lying in the outermost 4% of the volume of the detector; they include the events number 3,4,10 of the dataset, close to the threshold. We expect 1 background event in Baksan in  $T = 30$  s; this, together with a typical expected signal of about 1.6 antineutrinos ( $\mu = 2.6$ ) gives a reasonable Poisson chance  $\mu^n e^{-\mu}/n! = 8\%$  when  $n = 5$ . IMB instead can be assumed to be background free.

*Energy distribution* This distribution is difficult to interpret. We select 3 questions for the discussion:

- (i) Do the observed energies meet the expectations?
  - (ii) Are the average energies of IMB  $31.9 \pm 2.6$  MeV and of Kamiokande-II  $15.4 \pm 2.4$  MeV compatible?
  - (iii) Why the first 4 Kamiokande-II events have average energy  $12.6 \pm 2.4$  MeV and the last seven  $17.1 \pm 3.3$  MeV?
- The first can be answered fitting the data with a black body (thermal)  $\bar{\nu}_e$  spectrum with luminosity  $\propto R_c^2 T_c^4$ . We confirm that the Kamiokande-II and IMB datasets are compatible only when we consider the 90% C.L.; Baksan is compatible with both. When we combine the data in a unique analysis, the values of the radius is larger than expected,  $R_c \sim 30$  km. The second question can be discussed quantitatively calculating the average energies of the events, keeping the antineutrino temperature  $T_c$  as a free parameter (see e.g., [11]): Kamiokande-II and IMB suggest different values of  $T_c$ . The first two questions are the traditional formulations of the “energy problem”. They suggest two possible ways-out: a) The energy distribution is strongly non-thermal. This is explored, e.g., in [12] and [13]. Our objection is that this is a major deviation from the expectations that should not be admitted in a conservative analysis, as, e.g., [14]. b) Another possibility is that there are some low energy background events in Kamiokande-II [15], see the feature at  $E = 4 - 8$  MeV in Fig. 1, but recall that the background events above the nominal threshold are expected to be few. The third formulation of the problem [17],[15] offers a new clue: the excess of low energy Kamiokande-II events is due to the early detected events. This is fine




FIG. 1: Dots: observed energies in Kamiokande-II in 30 s. Continuous curve: expected cumulative spectrum (background + signal as in Eq. 3) as a function of the energy in MeV. The Smirnov-Cramèr-Von Mises (SCVM) [16] goodness of fit is 19%; compare also with [15].

since the standard scenario predicts the existence of the initial phase of accretion possibly with peculiar features. The high luminosity helps account the early events seen by Kamiokande-II.

*Angular distribution* The data and the expected angular distribution have been recently compared in [11], keeping into account the angular bias of IMB [5]. The SCVM test shows that the “problem” of the forward peak is not severe: if all event are IBD, the goodness of fit is 6.4% for IMB and 8.6% in Kamiokande-II. With 1 or 2 elastic scattering events (0.3-0.6 expected in Kamiokande-II) this improves further. Note that, even




FIG. 2: Expected 2-dimensional (positron-energy in MeV, cosine with the direction of SN1987A)-distribution for Kamiokande-II. The contour plot with equally spaced heights shows the mild directionality of IBD reaction and the peak of ES events (forward region near 15 MeV) whose shape is dictated by instrumental effects (see e.g., [11]).




FIG. 3: In the abscissa, the time of the event; in the ordinate, its progressive number. Two time distributions, comprising 22 signal and 7 background events, are given. Also indicated their goodness of fit values. In the lower one, all signal events belong to cooling, where the signal declines with a time constant  $\tau_c = 4$  s. In the upper one, only 13 signal events belong to cooling; the remaining 9 belong to accretion phase and are distributed in the interval  $t < 0.7$  s. The dotted vertical line marks the end of the first second.

being ready to consider something exotic, it is hard to imagine a reaction that is forward-peaked but too much, as needed to locate half of the IMB events in the region  $30^\circ < \theta < 60^\circ$ . Thus the discrepancy with the expectations is not very compelling. We also estimated the angular distribution taking into account elastic scattering events (ES). Assuming oscillations with normal mass hierarchy and increasing non-electronic neutrinos as allowed in [18] we calculated the time-averaged distribution of Fig. 2. From this, we find that the probability that the first event detected by Kamiokande-II is due to ES is about 30%.

*Time distribution* The time sequence of the Kamiokande-II events is at first sight odd; there is a cluster of six events in the first second and a “gap” of 7 seconds between the event number 9 and the event number 10, difficult to understand on physical basis. One way-out is to combine all data in a single dataset as in Fig. 3. This synthetic dataset can be compared with the expectations. Besides the known background component,  $n_b = 6.6$  expected events, the signal should accumulate in two phases: a first one with  $n_a$  events, followed by a second and longer phase with  $n_c = 29 - n_b - n_a$  events:

$$N(t) = n_b \frac{t}{T} + n_a [1 + \theta(t - \tau_a) (\frac{t}{\tau_a} - 1)] + n_c [1 - e^{-t/\tau_c}] \quad (1)$$

where the suffixes  $b, a, c$  mean background, accretion, cooling. The goodness-of-fit values shown in Fig. 3 suggest that a two component model of this type has no




FIG. 4: Continuous curve: instantaneous  $\bar{\nu}_e$  flux for  $M_a = 0.15 M_\odot$  and  $T_a = 2.5$  MeV. Dotted curve: black body distribution with the same luminosity ( $1.1 \times 10^{53}$  erg/s) and average energy (13 MeV), i.e., with parameters  $R_c = 82$  km and  $T_c = 4.1$  MeV.

problem to reproduce the data.<sup>1</sup> See [20, 21] for further useful discussion of the time sequence of the events.

### III. A TWO COMPONENT MODEL FOR $\bar{\nu}_e$ EMISSION

Let us summarize the previous section. There is some disturbing feature of the data. The presence of few elastic scattering events does not seem to be essential to explain the angular distribution, that comforts us to proceed with the usual IBD hypothesis for the signal. We showed that background events should be accounted for and that it is important to consider very seriously the existence of an initial phase of high neutrino luminosity. In short, the opinion that we derive from the previous discussion is that there is no serious problem with the data. Thus we move to interpret them within the ‘standard scenario’.

We analyzed the data of Kamiokande-II, IMB and Baksan adopting a definite model for neutrino emission [19]. The first attempt in this direction has been done by Lamb and Loredó [17]. Of course, the choice of the model is a critical step. We assume the existence of the two main phases of antineutrino emission as expected in the ‘standard scenario’. For each phase we need to describe the




FIG. 5: The continuous curve is the  $\bar{\nu}_e$  spectrum in the best fit model; the dotted curve is the approximant described in Eq. 3.

luminosity of the emission, the average antineutrino energy and the duration. For **cooling**, the parameters are  $R_c$ ,  $T_c$  and  $\tau_c$  (similar to the parameters of the previous section). For **accretion**, we use the mass of neutrons exposed to the positrons,  $M_a$ , the temperature of the positrons,  $T_a$ , and the duration of the accretion,  $\tau_a$ . In fact, the  $\bar{\nu}_e$  are produced from the reaction of the thermal positrons with the target neutrons around the proto neutron star, through  $e^+n \rightarrow p\bar{\nu}_e$ , the inverse of IBD; a sample energy spectrum is shown in Fig. 4. Note the characteristic ‘pinching’ of the distribution—an output, not an input of our model.

Let us mention for completeness some qualifying feature of our analysis. We prescribe that (1) the temperature of  $e^+$  increases during accretion so that the average energy of  $\bar{\nu}_e$  is approximatively continuous at  $t \sim \tau_a$ , that overcomes the shortcoming of the Lamb and Loredó model noted in [12]; (2) the number of neutrons exposed to the positron flux decreases in time more smoothly than as in [17]; in this way the luminosity is also continuous, as expected on general basis. (3) Finally we avoid the simultaneous presence of cooling and accretion  $\bar{\nu}_e$  delaying the cooling phase by an amount  $\sim \tau_a$  again improving on [17]. We also improved the energy spectrum of accretion neutrinos and included neutrino oscillations with normal mass hierarchy.

Our best fit result for the parameters is [19]:

$$\begin{aligned} R_c &= 16 \text{ km}, & T_c &= 4.6 \text{ MeV}, & \tau_c &= 4.7 \text{ s}, \\ M_a &= 0.22 M_\odot, & T_a &= 2.4 \text{ MeV}, & \tau_a &= 0.6 \text{ s}. \end{aligned} \quad (2)$$

The radiated binding energy is  $\mathcal{E}_b = 2.2 \times 10^{53}$  erg, similar to the expected value for the formation of a neutron star. The values of the parameters of the cooling phase are reasonable; in particular,  $R_c$  resembles a typical neutron star radius. Also the values of the parameters of accretion

<sup>1</sup> This still leaves some open questions: The absolute times of Kamiokande-II and Baksan are not reliable. Is it fair to use the corresponding freedom of interpretation to set the first event in each detectors at  $t = 0$ ? Do the assumed number of events fits into a reasonable model for the emission? Both of them are answered affirmatively with the global fit of the data discussed later, based on [19].

| $R(0)$ | $\beta$ | $N_{tot}$ | $\langle z \rangle$ | $N_{thr}$ | $\langle z \rangle$ | model    |
|--------|---------|-----------|---------------------|-----------|---------------------|----------|
| 2.00   | 2.00    | 3         | 0.5                 | 0.6       | 0.2                 | ref.[23] |
| 0.67   | 3.44    | 2         | 0.7                 | 0.3       | 0.2                 | ref.[13] |
| 1.25   | 3.44    | 4         | 0.7                 | 0.5       | 0.2                 | ref.[24] |

TABLE I: 1<sup>st</sup> and 2<sup>nd</sup> column, parameters of SN distribution— $R(0)$  is in units of  $1/(10^4 \text{ Mpc}^3 \text{ yr})$ ; 3<sup>rd</sup> and 4<sup>th</sup> column, total number of expected events per year from the relic supernovae and average redshift; 5<sup>th</sup> and 6<sup>th</sup> column, events above a threshold of 19.3 MeV and average redshift.

resemble the expectations; the value of initial accreting mass  $M_a$  is a fraction of the outer core mass  $\sim 0.6M_\odot$ . A last interesting outcome is that in the 30 s window Baksan had 0, 1, 2 background events with a posteriori probabilities of 20%, 47%, 29%, that compares well with the *a priori* expectation of 1 event.

The electron antineutrino spectrum, shown in Fig. 5 can be calculated from the fluence (=time integrated flux) assuming  $D = 50$  kpc, and it is well approximated by a modified Fermi-Dirac distribution:

$$\frac{dN}{dE} = \frac{\kappa}{T^3} \frac{E^2}{1 + \exp\left(\frac{E}{T} - \eta\right)} \quad \text{with} \quad \begin{cases} \kappa = 9.45 \times 10^{56}, \\ T = 3.77 \text{ MeV}, \\ \eta = 0.531. \end{cases} \quad (3)$$

The *a posteriori* comparison with the detected energies is satisfactory, as already shown in Fig. 1.

As a possible application, we use this result to predict the number of relic neutrino events. We consider a detector a la SK with a fiducial mass of 22.5 kton [22]. The rate of accumulation of IBD events is:

$$N_p \int_{19.3 \text{ MeV}}^{E_{max}} dE \sigma(E) \frac{c}{H_0} \int_0^{z_{max}} dz \frac{R(z) \frac{dN}{dE}((1+z)E)}{\sqrt{\Omega_\Lambda + \Omega_{DM}(1+z)^3}} \quad (4)$$

The cosmological parameters are  $H_0 = 70$  km/sec/Mpc,  $\Omega_\Lambda = 0.7$ ,  $\Omega_{DM} = 0.3$ . An important quantity is the rate of core collapse supernovae  $R$  as a function of the redshift  $z$ , adequately described by:  $R(z) = R(0)(1+z)^\beta$  if  $z < 1$  and  $= R(0)2^\beta$  if  $z \geq 1$ .  $R(0)$  is the product of the fraction of core collapse supernovae  $f_{SN}$  and the cosmic rate of star formation  $R_*(0)$ . Other quantities (the maximum energy  $E_\nu \sim 40 - 60$  MeV and redshift  $\sim 5 - 6$ , the point  $z = 1$  where the slope is modified) are not critical for the prediction. We compare in Tab. I the events per year for three descriptions of the rate of cosmic supernovae. The lower number of events is rather close to the one given in [13]; in other words, the non-standard neutrino spectrum used in [13] gives essentially the same results as the (more standard) spectrum of Eq. 3. Barring the possibility that SN1987A was a peculiar event, we are lead to believe that the largest uncertainty in predicting the relic neutrino signal is the rate of cosmic supernovae rather than the model for neutrino emission.

## IV. DISCUSSION

The difficulty to interpret SN1987A data is due to small-number statistics, partial operativity of IMB, poor absolute times in Kamiokande-II and Baksan, an apparent excess of events in Baksan, “gap” in the time-distribution of Kamiokande-II, peculiar angular distributions and different energies of Kamiokande-II and IMB. However, none of these problems constitutes an unsurmountable difficulty and on the contrary they can and have to be taken care in the analysis; furthermore, none of them can be solved too directly by considering reasonable deviations from the standard expectations. This does not guarantee us the possibility of reaching a safe interpretation of the data, but contributes to make less arbitrary the belief that these unique data can be understood in a coherent framework.

A general question that we addressed is if the data resemble the conventional expectations. We motivated the opinion that the presence of background events should be accounted for and that the signal should be described by considering an initial phase of intense neutrino luminosity, as required in the ‘standard scenario’ [8, 9]. This opinion is in agreement with the conclusions reached in 1989 in the review on SN1987A by Imshennik and Nadyozhin [25].

Being aware of the incompleteness of the present information (on 3D effects, rotation, magnetic fields; on peculiarities of SN1987A; on the existence of the neutron star; on neutrino oscillations; on the detailed detector response, *etc* [1]) we attempted to define the expectations of the ‘standard scenario’ [8, 9] in a parameterized model, trying to keep it as simple and flexible as possible but imposing certain general requirements, such as the continuity of the antineutrino luminosity and average energy.

We elaborated on the pioneer approach of Lamb and Loredó to SN1987A data analysis [17] emphasizing its defects and drawbacks but arguing that their model for antineutrino emission can be improved to resemble more closely the expectations [19]. The numerical analysis of the data yielded indications in (perhaps surprising) agreement with the expectations of the ‘standard scenario’ for neutrino emission. We estimated in this way the spectrum of  $\bar{\nu}_e$  from the SN1987A data and applied this result to evaluate the number of relic neutrinos events.

*Acknowledgments* F.V. thanks the Organizers of the Symposium for the honorable invitation; O. Ryazhskaya for many stimulating discussions and inexhaustible passion with SN1987A; D. Nadyozhin for help with astrophysics and precious encouragement; E. Alexeev and A. Mal’gin for frank and useful discussions on the meaning of the data; and last but not least my friends N. Agafonova, A. Mal’gin, and V. Yakushev who made my first visit to Moscow unforgettable.

- 
- [1] Web page: <http://vaxmw.tower.ras.ru/SN1987A>
  - [2] V. L. Dadykin *et al.*, JETP Lett. **45** (1987) 593 [Pisma Zh. Eksp. Teor. Fiz. **45** (1987) 464]
  - [3] V.S. Imshennik, O.G. Ryazhskaya, Astron. Lett. **30** (2004) 14
  - [4] K. Hirata *et al.* [KAMIOKAND-II Collaboration], Phys. Rev. Lett. **58** (1987) 1490; Phys. Rev. D **38**, 448-458 (1988)
  - [5] R.M. Bionta *et al.*, Phys. Rev. Lett. **58** (1987) 1494; C.B. Bratton *et al.*, IMB collaboration, Phys. Rev. D **37** (1988) 3361
  - [6] E.N. Alekseev, L.N. Alekseeva, V.I. Volchenko and I.V. Krivosheina, JETP Lett. **45** (1987) 589 and Phys.Lett. **B 205** (1988) 209
  - [7] V.L. Dadykin, G.T. Zatsepin, and O.G. Ryazhskaya, Sov. Phys. Usp., **32** (5) 1989; O.G. Ryazhskaya, Usp. Fiz. Nauk **49** (10) 2006
  - [8] D.K. Nadyozhin, Astrophysics & Space Science **59** (1978) 131
  - [9] J.R. Wilson in *Numerical Astrophysics*, eds. J. Centrella, J. LeBlanc, R.L. Bowers, page 422 (Jones and Bartlett, Boston, 1985); H.A. Bethe and J.R. Wilson, Astrophys. J. **295** (1985) 14
  - [10] H.-T. Janka *et al.*, 0708.3372 [astro-ph]
  - [11] M.L. Costantini *et al.*, Phys.Rev.D**70** (2004) 043006
  - [12] A. Mirizzi and G. G. Raffelt, Phys. Rev. D **72** (2005) 063001
  - [13] C. Lunardini, Astropart. Ph. **26** (2006) 190
  - [14] H.-T. Janka and W. Hillebrandt, A& A **224** (1989) 49
  - [15] M.L. Costantini, A. Ianni, G. Pagliaroli, F. Vissani, JCAP **0708:014**, 2007.
  - [16] Obtained calculating  $\omega^2 = 1/12n + \sum_{i=1}^n (F(E_i) - (i-1/2)/n)^2$ , where  $n$ =number of data and  $F(E)$ =cumulative distribution
  - [17] T.J. Loredo and D.Q. Lamb, Phys. Rev. D **65**, 063002, 2002
  - [18] M.T. Keil *et al.*, Astrophys.J. **590** (2003) 971
  - [19] G. Pagliaroli, M.L. Costantini, A. Ianni, F. Vissani, LNGS/TH-01/08, For a preliminary report, see 0708.4032 [astro-ph]
  - [20] A. Mal'gin, Nuovo Cim. C **21** (1998) 317.
  - [21] V.S. Imshennik, D.K. Nadyozhin, IJMP A**20** (2005) 6597
  - [22] M. Malek *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **90** (2003) 061101
  - [23] L.E. Strigari *et al.*, JCAP **0504:017**, 2005
  - [24] H. Yuksel and J.F. Beacom, Phys. Rev. D **76** (2007) 083007
  - [25] V.S. Imshennik, D.K. Nadyozhin, Astrophysics & Space Science Physics Review **8** (1989) part 1